

Redox

- [Redox: From Wikipedia, the free encyclopedia](#)
- [Redox in a Water Fuel Cell](#)

Redox: From Wikipedia, the free encyclopedia


Source: <https://en.wikipedia.org/wiki/Redox>

From Wikipedia, the free encyclopedia

[Jump to navigation](#)[Jump to search](#)

For other uses, see [Redox \(disambiguation\)](#).

Image not found or type unknown




Sodium and fluorine bonding ionically to form sodium fluoride. Sodium loses its outer electron to give it a stable electron configuration, and this electron enters the fluorine atom exothermically. The oppositely charged ions are then attracted to each other. The sodium is oxidized; and the fluorine is reduced.

0:47

Demonstration of the reaction between a strong oxidizing and a reducing agent. When a few drops of glycerol (mild reducing agent) are added to powdered potassium permanganate (strong oxidizing agent), a violent redox reaction accompanied by self-ignition starts.

Image not found or type unknown



Example of a reduction–oxidation reaction between sodium and chlorine, with the *OIL RIG* mnemonic^[1]

Redox (reduction-oxidation, /'rɛdɒks/ RED-oks, /'ri:dɒks/ REE-doks^[2]) is a type of chemical reaction in which the oxidation states of substrate change.^[3]

- **Oxidation** is the *loss* of electrons or an *increase* in the oxidation state of a chemical or atoms within it.
- **Reduction** is the *gain* of electrons or a *decrease* in the oxidation state of a chemical or atoms within it.

There are two classes of redox reactions:

- *electron-transfer* where only one (usually) electron flows from the reducing agent to the oxidant. This type of redox reaction is often discussed in term of redox couples and electrode potentials.
- *atom transfer*, where an atom transfers from one substrate to another. For example, in the rusting of iron, the oxidation state of iron atoms increases as it converts to an oxide and simultaneously the oxidation state of oxygen decreases as it accepts electrons released by iron. Although oxidation reactions are commonly associated with the formation of oxides, other chemical species can serve the same function.^[4] In hydrogenation, C=C (and other) bonds are reduced by transfer of hydrogen atoms.

□

Contents

- 1Terminology
 - 1.1Oxidants
 - 1.2Reducers
 - 1.3Electronation and deelectronation
- 2Rates, mechanisms, and energies
- 3Standard electrode potentials (reduction potentials)
- 4Examples of redox reactions
 - 4.1Metal displacement

- 4.2 Other examples
- 4.3 Corrosion and rusting
- 4.4 Disproportionation
- 5 Redox reactions in industry
- 6 Redox reactions in biology
 - 6.1 Redox cycling
- 7 Redox reactions in geology
- 8 Redox reactions in soils
- 9 Mnemonics
- 10 See also
- 11 References
- 12 Further reading
- 13 External links

Terminology

"Redox" is a combination of the words "reduction" and "oxidation". The term "redox" was first used in 1928.^[5] The processes of oxidation and reduction occur simultaneously and cannot occur independently.^[4] In redox processes, the reductant transfers electrons to the oxidant. Thus, in the reaction, the reductant or *reducing agent* loses electrons and is oxidized, and the oxidant or *oxidizing agent* gains electrons and is reduced. The pair of an oxidizing and reducing agent that is involved in a particular reaction is called a *redox pair*. A *redox couple* is a reducing species and its corresponding oxidizing form,^[6] e.g., Fe^{2+}

/ Fe^{3+}

.The oxidation alone and the reduction alone are each called a *half-reaction* because two half-reactions always occur together to form a whole reaction.

Oxidants


Oxidation originally implied reaction with oxygen to form an oxide. Later, the term was expanded to encompass oxygen-like substances that accomplished parallel chemical reactions. Ultimately, the meaning was generalized to include all processes involving the loss of electrons. Substances that have the ability to *oxidize* other substances (cause them to lose electrons) are said to be *oxidative* or *oxidizing*, and are known as *oxidizing agents*, *oxidants*, or *oxidizers*. The oxidant (oxidizing agent) removes electrons from another substance, and is thus itself reduced. And,

because it "accepts" electrons, the oxidizing agent is also called an electron acceptor. Oxidants are usually chemical substances with elements in high oxidation states (e.g., H

2^0
 $2, \text{MnO}^-$
 $4, \text{CrO}$
 $3, \text{Cr}$
 2O^{2-}
 $7, \text{OsO}$

4), or else highly electronegative elements (O_2 , F_2 , Cl_2 , Br_2) that can gain extra electrons by oxidizing another substance.^[*citation needed*]

Oxidizers are oxidants but the term is mainly reserved for sources of oxygen, particularly in the context of explosions. Nitric acid is an oxidizer.



The international pictogram for oxidizing chemicals

Main article: Oxidizing agent

Oxygen is the quintessential oxidizer.

Reducers

Main article: Reducing agent

Substances that have the ability to *reduce* other substances (cause them to gain electrons) are said to be *reductive* or *reducing* and are known as *reducing agents*, reductants, or reducers. The reductant (reducing agent) transfers electrons to another substance and is thus itself oxidized. And, because it donates electrons, the reducing agent is also called an electron donor. Electron donors can also form charge transfer complexes with electron acceptors. The word *reduction* originally referred to the loss in weight upon heating a metallic ore such as a metal oxide to extract the metal. In other words, ore was "reduced" to metal. Antoine Lavoisier demonstrated that this loss of weight was due to the loss of oxygen as a gas. Later, scientists realized that the metal

atom gains electrons in this process. The meaning of *reduction* then became generalized to include all processes involving a gain of electrons. **Reducing equivalent** refers to chemical species which transfer the equivalent of one electron in redox reactions. The term is common in biochemistry.[7] A reducing equivalent can be an electron, a hydrogen atom, as a hydride ion.[8]

Reductants in chemistry are very diverse. Electropositive elemental metals, such as lithium, sodium, magnesium, iron, zinc, and aluminium, are good reducing agents. These metals donate or *give away* electrons relatively readily. They transfer electrons.

Hydride transfer reagents, such as NaBH_4 and LiAlH_4 , reduce by atom transfer: they transfer the equivalent of hydride or H^- . These reagents widely used in [the reduction of carbonyl compounds to alcohols.[9][10] A related method of reduction involves the use of hydrogen gas (H_2) as sources of H atoms.

Electronation and deelectronation

The electrochemist John Bockris proposed the words *electronation* and *deelectronation* to describe reduction and oxidation processes, respectively, when they occur at electrodes.[11] These words are analogous to protonation and deprotonation.[12] They have not been widely adopted by chemists worldwide, although IUPAC has recognized the term electronation.[13]

Rates, mechanisms, and energies


Redox reactions can occur slowly, as in the formation of rust, or rapidly, as in the case of burning fuel. Electron transfer reactions are generally fast, occurring within the time of mixing.

The mechanisms of atom-transfer reactions are highly variable because many kinds of atoms can be transferred. Such reactions can also be quite complex, i.e. involve many steps. The mechanisms of electron-transfer reactions occur by two distinct pathways, inner sphere electron transfer and outer sphere electron transfer.

Analysis of bond energies and ionization energies in water allow calculation of the thermodynamic aspects of redox reactions.

Standard electrode potentials (reduction potentials)

Each half-reaction has a *standard electrode potential* (E°_{cell}), which is equal to the potential difference or voltage at equilibrium under standard conditions of an electrochemical cell in which the cathode reaction is the half-reaction considered, and the anode is a standard hydrogen electrode where hydrogen is oxidized:



The electrode potential of each half-reaction is also known as its *reduction potential* E°_{red} , or potential when the half-reaction takes place at a cathode. The reduction potential is a measure of the tendency of the oxidizing agent to be reduced. Its value is zero for $\text{H}^+ + \text{e}^- \rightarrow \frac{1}{2} \text{H}_2$ by definition, positive for oxidizing agents stronger than H^+ (e.g., +2.866 V for F_2) and negative for oxidizing agents that are weaker than H^+ (e.g., -0.763 V for Zn^{2+}).^[14]

For a redox reaction that takes place in a cell, the potential difference is:

$$E^\circ_{\text{cell}} = E^\circ_{\text{cathode}} - E^\circ_{\text{anode}}$$

However, the potential of the reaction at the anode is sometimes expressed as an *oxidation potential*:

$$E^\circ_{\text{ox}} = -E^\circ_{\text{red}}$$

The oxidation potential is a measure of the tendency of the reducing agent to be oxidized but does not represent the physical potential at an electrode. With this notation, the cell voltage equation is written with a plus sign

$$E^\circ_{\text{cell}} = E^\circ_{\text{red(cathode)}} + E^\circ_{\text{ox(anode)}}$$

Examples of redox reactions






Illustration of a redox reaction


In the reaction between hydrogen and fluorine, hydrogen is being oxidized and fluorine is being reduced:



This reaction is spontaneous and releases 542 kJ per 2 g of hydrogen because the H-F bond is much stronger than the F-F bond. This reaction can be analyzed as two half-reactions. The oxidation reaction converts hydrogen to protons:



The reduction reaction converts fluorine to the fluoride anion:




The half reactions are combined so that the electrons cancel:

H_2	\rightarrow	$2 \text{H}^+ + 2 \text{e}^-$
$\text{F}_2 + 2 \text{e}^-$	\rightarrow	2F^-
<hr/>		
$\text{H}_2 + \text{F}_2$	\rightarrow	$2 \text{H}^+ + 2 \text{F}^-$

The protons and fluoride combine to form hydrogen fluoride in a non-redox reaction:




The overall reaction is:




Metal displacement

Image not found or type unknown




A redox reaction is the force behind an electrochemical cell like the Galvanic cell pictured. The battery is made out of a zinc electrode in a ZnSO_4 solution connected with a wire and a porous disk to a copper electrode in a CuSO_4 solution.

In this type of reaction, a metal atom in a compound (or in a solution) is replaced by an atom of another metal. For example, copper is deposited when zinc metal is placed in a copper(II) sulfate solution:




In the above reaction, zinc metal displaces the copper(II) ion from copper sulfate solution and thus liberates free copper metal. The reaction is spontaneous and releases 213 kJ per 65 g of zinc.

The ionic equation for this reaction is:



As two half-reactions, it is seen that the zinc is oxidized:



And the copper is reduced:



Other examples


- The reduction of nitrate to nitrogen in the presence of an acid (denitrification):
$$\{ \text{2NO}_3^- + 10\text{e}^- + 12\text{H}^+ \rightarrow \text{N}_2 + 6 \text{H}_2\text{O} \}$$

$$\{ \text{2NO}_3^- + 10\text{e}^- + 12\text{H}^+ \rightarrow \text{N}_2 + 6 \text{H}_2\text{O} \}$$
- The combustion of hydrocarbons, such as in an internal combustion engine, produces water, carbon dioxide, some partially oxidized forms such as carbon monoxide, and heat energy. Complete oxidation of materials containing carbon produces carbon dioxide.
- The stepwise oxidation of a hydrocarbon by oxygen, in organic chemistry, produces water and, successively: an alcohol, an aldehyde or a ketone, a carboxylic acid, and then a peroxide.

Corrosion and rusting



Oxides, such as iron(III) oxide or rust, which consists of hydrated iron(III) oxides $\text{Fe}_2\text{O}_3 \cdot n\text{H}_2\text{O}$ and iron(III) oxide-hydroxide ($\text{FeO}(\text{OH})$, $\text{Fe}(\text{OH})_3$), form when oxygen combines with other elements






Iron rusting in pyrite cubes



- The term **corrosion** refers to the electrochemical oxidation of metals in reaction with an oxidant such as oxygen. **Rusting**, the formation of iron oxides, is a well-known example of electrochemical corrosion; it forms as a result of the oxidation of iron metal. Common rust often refers to iron(III) oxide, formed in the following chemical reaction:



- The oxidation of iron(II) to iron(III) by hydrogen peroxide in the presence of an acid:




Here the overall equation involves adding the reduction equation to twice the oxidation equation, so that the electrons cancel:



Disproportionation

A disproportionation reaction is one in which a single substance is both oxidized and reduced. For example, thiosulfate ion with sulfur in oxidation state +2 can react in the presence of acid to form elemental sulfur (oxidation state 0) and sulfur dioxide (oxidation state +4).



Thus one sulfur atom is reduced from +2 to 0, while the other is oxidized from +2 to +4.[15]

Redox reactions in industry

Cathodic protection is a technique used to control the corrosion of a metal surface by making it the cathode of an electrochemical cell. A simple method of protection connects protected metal to a more easily corroded "sacrificial anode" to act as the anode. The sacrificial metal instead of the protected metal, then, corrodes. A common application of cathodic protection is in galvanized steel, in which a sacrificial coating of zinc on steel parts protects them from rust.^[citation needed]

Oxidation is used in a wide variety of industries such as in the production of cleaning products and oxidizing ammonia to produce nitric acid.

Redox reactions are the foundation of electrochemical cells, which can generate electrical energy or support electrosynthesis. Metal ores often contain metals in oxidized states such as oxides or sulfides, from which the pure metals are extracted by smelting at high temperature in the presence of a reducing agent. The process of electroplating uses redox reactions to coat objects with a thin layer of a material, as in chrome-plated automotive parts, silver plating cutlery, galvanization and gold-plated jewelry.^[citation needed]

Redox reactions in biology

ascorbic acid


Image not found or type unknown

dehydroascorbic acid

Image not found or type unknown

Top: ascorbic acid (reduced form of Vitamin C)


Bottom: dehydroascorbic acid (oxidized form of Vitamin C)



Enzymatic browning is an example of a redox reaction that takes place in most fruits and vegetables.

Many important biological processes involve redox reactions. Before some of these processes can begin iron must be assimilated from the environment.[16]

Cellular respiration, for instance, is the oxidation of glucose ($C_6H_{12}O_6$) to CO_2 and the reduction of oxygen to water. The summary equation for cell respiration is:



The process of cell respiration also depends heavily on the reduction of NAD^+ to NADH and the reverse reaction (the oxidation of NADH to NAD^+). Photosynthesis and cellular respiration are complementary, but photosynthesis is not the reverse of the redox reaction in cell respiration:



Biological energy is frequently stored and released by means of redox reactions. Photosynthesis involves the reduction of carbon dioxide into sugars and the oxidation of water into molecular oxygen. The reverse reaction, respiration, oxidizes sugars to produce carbon dioxide and water. As intermediate steps, the reduced carbon compounds are used to reduce nicotinamide adenine dinucleotide (NAD^+) to NADH, which then contributes to the creation of a proton gradient, which drives the synthesis of adenosine triphosphate (ATP) and is maintained by the reduction of oxygen. In animal cells, mitochondria perform similar functions. See the *Membrane potential* article.


Free radical reactions are redox reactions that occur as a part of homeostasis and killing microorganisms, where an electron detaches from a molecule and then reattaches almost instantaneously. Free radicals are a part of redox molecules and can become harmful to the human body if they do not reattach to the redox molecule or an antioxidant. Unsatisfied free radicals can spur the mutation of cells they encounter and are, thus, causes of cancer.

The term *redox state* is often used to describe the balance of GSH/GSSG, NAD⁺/NADH and NADP⁺/NADPH in a biological system such as a cell or organ. The redox state is reflected in the balance of several sets of metabolites (e.g., lactate and pyruvate, beta-hydroxybutyrate, and acetoacetate), whose interconversion is dependent on these ratios. An abnormal redox state can develop in a variety of deleterious situations, such as hypoxia, shock, and sepsis. Redox mechanism also control some cellular processes. Redox proteins and their genes must be co-located for redox regulation according to the CoRR hypothesis for the function of DNA in mitochondria and chloroplasts.

Redox cycling

Wide varieties of aromatic compounds are enzymatically reduced to form free radicals that contain one more electron than their parent compounds. In general, the electron donor is any of a wide variety of flavoenzymes and their coenzymes. Once formed, these anion free radicals reduce molecular oxygen to superoxide and regenerate the unchanged parent compound. The net reaction is the oxidation of the flavoenzyme's coenzymes and the reduction of molecular oxygen to form superoxide. This catalytic behavior has been described as a *futile cycle* or redox cycling.

Redox reactions in geology




Blast furnaces of Třinec Iron and Steel Works, Czech Republic

Minerals are generally oxidized derivatives of metals. Iron is mined as its magnetite (Fe_3O_4).

Titanium is mined as its dioxide, usually in the form of rutile (TiO_2). To obtain the corresponding metals, these oxides must be reduced, which is often achieved by heating these oxides with carbon or carbon monoxide as reducing agents. Blast furnaces are the reactors where iron oxides and coke (a form of carbon) are combined to produce molten iron. The main chemical reaction producing the

molten iron is:[17]



Redox reactions in soils[edit]

Electron transfer reactions are central to myriad processes and properties in soils, and electron "activity", quantified as Eh (platinum electrode potential (voltage) relative to the standard hydrogen electrode) or pe (analogous to pH as -log electron activity), is a master variable, along with pH, that controls and is governed by chemical reactions and biological processes. Early theoretical research with applications to flooded soils and paddy rice production was seminal for subsequent work on thermodynamic aspects of redox and plant root growth in soils.[18] Later work built on this foundation, and expanded it for understanding redox reactions related to heavy metal oxidation state changes, pedogenesis and morphology, organic compound degradation and formation, free radical chemistry, wetland delineation, soil remediation, and various methodological approaches for characterizing the redox status of soils.[19][20]

Mnemonics[edit]

Main article: [List of chemistry mnemonics](#)

The key terms involved in redox can be confusing.[21][22] For example, a reagent that is oxidized loses electrons; however, that reagent is referred to as the reducing agent. Likewise, a reagent that is reduced gains electrons and is referred to as the oxidizing agent.[23] These mnemonics are commonly used by students to help memorise the terminology:[24]

- "OIL RIG" — **o**xidation **i**s **l**oss of electrons, **r**eduction **i**s **g**ain of electrons[21][22][23][24]
- "LEO the lion says GER [grr]" — **l**oss of **e**lectrons is **o**xidation, **g**ain of **e**lectrons is **r**eduction[21][22][23][24]
- "LEORA says GEROA" — the loss of electrons is called oxidation (reducing agent); the gain of electrons is called reduction (oxidizing agent).[23]
- "RED CAT" and "AN OX", or "AnOx RedCat" ("an ox-red cat") — reduction occurs at the cathode and the anode is for oxidation
- "RED CAT gains what AN OX loses" - reduction at the cathode gains (electrons) what anode oxidation loses (electrons)

- "PANIC" - Positive Anode and Negative is Cathode. This applies to electrolytic cells which release stored electricity, and can be recharged with electricity. PANIC does not apply to cells that can be recharged with redox materials. These galvanic or voltaic cells, such as fuel cells, produce electricity from internal redox reactions. Here, the positive electrode is the cathode and the negative is the anode.

Redox in a Water Fuel Cell

Capacitors are devices used to store electrical energy. They consist of two conductive plates separated by a dielectric material. When a voltage is applied across the plates, an electric field is created, which causes the plates to store energy. Capacitors are commonly used in electronic circuits for filtering, timing, and energy storage.

A water-filled capacitor is a special type of capacitor in which one of the conductive plates is replaced with water. The water serves as the dielectric material, and the other plate is usually made of metal. When an electric field is applied to the water-filled capacitor, an interesting phenomenon occurs: a red material forms at the interface between the water and the metal plate.

The red material that forms in a water-filled capacitor is due to a redox reaction. Redox is short for reduction-oxidation, which is a chemical reaction in which one substance loses electrons (oxidation) and another substance gains electrons (reduction). In a water-filled capacitor, the metal plate serves as the oxidizing agent, and the water serves as the reducing agent.

The source of the red material that forms in a water-filled capacitor is the metal plate. When an electric field is applied across the plates of the capacitor, electrons flow from the negative plate to the positive plate. The metal plate, which is the positive plate, attracts electrons from the water, which is the negative plate. As electrons are transferred from the water to the metal plate, the water becomes positively charged and the metal plate becomes negatively charged.

The transfer of electrons from the water to the metal plate causes a redox reaction to occur. The metal plate oxidizes the water, causing the water to lose electrons. The electrons that are lost by the water are then transferred to the metal plate, causing it to reduce. The red material that forms at the interface between the water and the metal plate is the product of this redox reaction.

The red material that forms in a water-filled capacitor can be a variety of compounds, depending on the metal used in the capacitor. For example, if the metal plate is made of iron, the red material that forms is likely to be iron oxide or rust. If the metal plate is made of copper, the red material that forms is likely to be copper oxide.

In conclusion, the red material that forms in a water-filled capacitor when an electric field is applied is due to a redox reaction between the metal plate and the water. The metal plate acts as the oxidizing agent, and the water acts as the reducing agent. The red material that forms is the product of this reaction, and its composition depends on the metal used in the capacitor. While the formation of the red material may be interesting to observe, it can also have a negative impact on the performance of the capacitor, as it can reduce its capacitance and increase its resistance.